

Datasheet for ABIN7127268

Recombinant anti-HIST1H3A antibody (H3K14ac)[Go to Product page](#)**3** Images

Overview

Quantity:	100 µL
Target:	HIST1H3A
Binding Specificity:	H3K14ac
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This HIST1H3A antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide
Clone:	3H3
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Purification:	Affinity-chromatography

Target Details

Target:	HIST1H3A
Alternative Name:	HIST1H3A (HIST1H3A Products)

Target Details


Background:	Background: Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. Aliases: Histone H3.1, Histone H3/a, Histone H3/b, Histone H3/c, Histone H3/d, Histone H3/f, Histone H3/h, Histone H3/i, Histone H3/j, Histone H3/k, Histone H3/l, HIST1H3A, H3FA, AND, HIST1H3B, H3FL, AND, HIST1H3C, H3FC, AND, HIST1H3D, H3FB, AND, HIST1H3E, H3FD, AND, HIST1H3F, H3FI, AND, HIST1H3G, H3FH, AND, HIST1H3H, H3FK, AND, HIST1H3I, H3FF, AND, HIST1H3J, H3FJ
UniProt:	P68431

Application Details

Application Notes:	Recommended dilution: WB:1:500-1:2000, ICC:1:50-1:500, IF:1:30-1:200,
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	Rabbit IgG in phosphate buffered saline, pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunofluorescence

Image 1. Immunofluorescence staining of HeLa cells (treated by 15 mM sodium butyrate for 30 min) with ABIN7127268 at 1:46, counter-stained with DAPI. The cells were fixed in 4 % formaldehyde, permeabilized using 0.2 % Triton X-100 and blocked in 10 % normal Goat Serum. The cells were then incubated with the antibody overnight at 4 °C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



Western Blotting

Image 2. Western Blot Positive WB detected in HeLa whole cell lysate, HepG2 whole cell lysate, SH-SY5Y whole cell lysate, and brain tissue. All lanes: Acetyl-Histone H3.1(K14) antibody at 0.75 µg/mL. Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 15 KDa. Observed band size: 15 KDa.



Immunocytochemistry

Image 3. Immunocytochemistry analysis of ABIN7127268 diluted at 1:100 and staining in HeLa cells performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.