
antibodies - online.com







Recombinant anti-HTR2C antibody

Images



\sim			
	$ \backslash / \square $	r\/I	\square

Quantity:	100 μL
Target:	HTR2C
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This HTR2C antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	A synthesized peptide derived from human 5HT2C Receptor	
Clone:	9H8	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Affinity-chromatography	

Target Details

Target:	HTR2C
Alternative Name:	HTR2C (HTR2C Products)
Background: Background: G-protein coupled receptor for 5-hydroxytryptamine (serotonin). Also function	

a receptor for various drugs and psychoactive substances, including ergot alkaloid derivatives, 1-2,5,-dimethoxy-4-iodophenyl-2-aminopropane (DOI) and lysergic acid diethylamide (LSD). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors. Beta-arrestin family members inhibit signaling via G proteins and mediate activation of alternative signaling pathways. Signaling activates a phosphatidylinositol-calcium second messenger system that modulates the activity of phosphatidylinositol 3-kinase and down-stream signaling cascades and promotes the release of Ca(2+) ions from intracellular stores. Regulates neuronal activity via the activation of short transient receptor potential calcium channels in the brain, and thereby modulates the activation of pro-opiomelacortin neurons and the release of CRH that then regulates the release of corticosterone. Plays a role in the regulation of appetite and eating behavior, responses to anxiogenic stimuli and stress. Plays a role in insulin sensitivity and glucose homeostasis.

Aliases: 5-hydroxytryptamine receptor 2C (5-HT-2C) (5-HT2C) (5-HTR2C) (5-hydroxytryptamine receptor 1C) (5-HT-1C) (5-HT1C) (Serotonin receptor 2C), HTR2C, HTR1C

UniProt:

P28335

Pathways:

Inositol Metabolic Process, Regulation of Carbohydrate Metabolic Process, Feeding Behaviour


Application Details


Application Notes: Recommended dilution: WB:1:500-1:5000, IHC:1:50-1:200,

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	Rabbit IgG in phosphate buffered saline, pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.





Immunohistochemistry

Image 1. IHC image of ABIN7127314 diluted at 1:100 and staining in paraffin-embedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05 % DAB.

Western Blotting

Image 2. Western Blot Positive WB detected in: Hela whole cell lysate, HepG2 whole cell lysate, A549 whole cell lysate All lanes: HTR2C antibody at 1:2000 Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 52, 29 kDa Observed band size: 75 kDa