

Datasheet for ABIN7127830

Recombinant anti-SUMO1 antibody[Go to Product page](#)**4** Images

Overview

| | |
|----------------|--|
| Quantity: | 100 µL |
| Target: | SUMO1 |
| Reactivity: | Human |
| Host: | Rabbit |
| Antibody Type: | Recombinant Antibody |
| Clonality: | Monoclonal |
| Conjugate: | This SUMO1 antibody is un-conjugated |
| Application: | ELISA, Immunohistochemistry (IHC), Flow Cytometry (FACS), Immunoprecipitation (IP) |

Product Details

| | |
|-------------------|--|
| Immunogen: | A synthesized peptide derived from human SUMO1 |
| Clone: | 5G3 |
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | Affinity-chromatography |

Target Details

| | |
|-------------------|--|
| Target: | SUMO1 |
| Alternative Name: | SUMO1 (SUMO1 Products) |
| Background: | Background: Ubiquitin-like protein that can be covalently attached to proteins as a monomer or |

Target Details

a lysine-linked polymer. Covalent attachment via an isopeptide bond to its substrates requires prior activation by the E1 complex SAE1-SAE2 and linkage to the E2 enzyme UBE2I, and can be promoted by E3 ligases such as PIAS1-4, RANBP2 or CBX4. This post-translational modification on lysine residues of proteins plays a crucial role in a number of cellular processes such as nuclear transport, DNA replication and repair, mitosis and signal transduction. Involved for instance in targeting RANGAP1 to the nuclear pore complex protein RANBP2. Covalently attached to the voltage-gated potassium channel KCNB1, this modulates the gating characteristics of KCNB1 (PubMed:19223394). Polymeric SUMO1 chains are also susceptible to polyubiquitination which functions as a signal for proteasomal degradation of modified proteins. May also regulate a network of genes involved in palate development. Covalently attached to ZFHX3 (PubMed:24651376).

Aliases: Small ubiquitin-related modifier 1, SUMO-1, GAP-modifying protein 1, GMP1, SMT3 homolog 3, Sentrin, Ubiquitin-homology domain protein PIC1, Ubiquitin-like protein SMT3C, Smt3C, Ubiquitin-like protein UBL1, SUMO1, SMT3C, SMT3H3, UBL1, OK/SW-cl.43

UniProt: [P63165](#)

Pathways: [M Phase](#), [Positive Regulation of Endopeptidase Activity](#), [Protein targeting to Nucleus](#), [Ubiquitin Proteasome Pathway](#)

Application Details

Application Notes: Recommended dilution: IHC:1:50-1:200, IP:1:200-1:1000,

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.




Immunohistochemistry

Image 1. IHC image of ABIN7127830 diluted at 1:92.5 and staining in paraffin-embedded human liver cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunohistochemistry

Image 2. IHC image of ABIN7127830 diluted at 1:92.5 and staining in paraffin-embedded human adrenal gland tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Western Blotting

Image 3. Immunoprecipitating SUMO1 in 293T whole cell lysate Lane 1: Rabbit control IgG instead of ABIN7127830 in 293T whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000) Lane 2: ABIN7127830 (3 µg) + 293T whole cell lysate (500 µg) Lane 3: 293T whole cell lysate (20 µg)

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN7127830.