

Datasheet for ABIN7129263

anti-DSTYK antibody[Go to Product page](#)**2** Images

Overview

Quantity:	100 µL
Target:	DSTYK
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DSTYK antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Fusion protein of Human DSTYK
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Purification:	Antigen affinity purification

Target Details

Target:	DSTYK
Alternative Name:	DSTYK (DSTYK Products)
Background:	Background: This gene encodes a dual serine/threonine and tyrosine protein kinase which is expressed in multiple tissues. It is thought to function as a regulator of cell death. Multiple transcript variants encoding different isoforms have been found for this gene. Aliases: DSTYK antibody, Dual serine/threonine and tyrosine protein kinase antibody, Dusty PK

Target Details

antibody, Dusty protein kinase antibody, DUSTY_HUMAN antibody, HDCMD38P antibody, KIAA0472 antibody, Receptor interacting protein 5 antibody, Receptor interacting protein kinase 5 antibody, Receptor interacting serine/threonine protein kinase 5 antibody, Receptor-interacting serine/threonine-protein kinase 5 antibody, RIP homologous kinase antibody, RIP-homologous kinase antibody, RIP5 antibody, RIPK5 antibody, SgK496 antibody, Sugem kinase 496 antibody

UniProt: [Q6XUX3](#)

Application Details

Application Notes: ELISA:1:1000-1:2000, WB:1:200-1:1000, IHC:1:25-1:100,

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: -20 °C, pH 7.4 PBS, 0.05 % Sodium azide, 40 % Glycerol


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using ABIN7129263(DSTYK Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original magnification: x200)



Western Blotting

Image 2. Gel: 6 % SDS-PAGE, Lysate: 40 µg, Lane 1-2: 823 cells, hepg2 cells, Primary antibody: ABIN7129263(DSTYK Antibody) at dilution 1/200, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 1 minute