
# antibodies - online.com











### Overview

Quantity:	100 μL
Target:	AXIN2
Binding Specificity:	AA 331-430
Reactivity:	Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This AXIN2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Axin 2
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Horse
Purification:	Purified by Protein A.
T	

## Target Details

Target: AXIN2

## **Target Details**


Alternative Name:	Axin 2 (AXIN2 Products)
Background:	Synonyms: AXIL, ODCRCS, Axin-2, Axin-like protein, Axis inhibition protein 2, Conductin, AXIN2 Background: Inhibitor of the Wnt signaling pathway. Down-regulates beta-catenin. Probably facilitate the phosphorylation of beta-catenin and APC by GSK3B (By similarity).
Gene ID:	8313
UniProt:	Q9Y2T1
Pathways:	Regulation of G-Protein Coupled Receptor Protein Signaling

# Application Details

Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only


# Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months



## **Immunohistochemistry (Paraffin-embedded Sections)**

**Image 1.** Formalin-fixed and paraffin embedded rat lymph node labeled with Rabbit Anti-Axin 2 Polyclonal Antibody, Unconjugated at 1:200 followed by conjugation to the secondary antibody and DAB staining



### **Immunohistochemistry**

**Image 2.** Formalin-fixed and paraffin embedded rat brain tissue labeled with Anti Axin 2 Polyclonal Antibody,Unconjugated (ABIN713696) at 1:200 followed by conjugation to the secondary antibody and DAB staining.