antibodies.com

Datasheet for ABIN7138433 anti-MYL9 antibody (pSer15)

2 Images



Overview

| Quantity: | 100 µL |
|----------------------|---|
| Target: | MYL9 |
| Binding Specificity: | pSer15 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Application: | Western Blotting (WB), ELISA, Immunofluorescence (IF) |

Product Details

| Immunogen: | Peptide sequence around phosphorylation site of serine 15 (A-N-S(p)-N-V) derived from Human MLC2. | |
|-------------------|---|--|
| Isotype: | lgG | |
| Cross-Reactivity: | Human | |
| Purification: | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi | |

Target Details

| Target: | MYL9 | |
|-------------------|----------------------|--|
| Alternative Name: | MYL9 (MYL9 Products) | |
| Background: | Background: | |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7138433 | 01/16/2024 | Copyright antibodies-online. All rights reserved.

| | Myosin regulatory subunit that plays an important role in regulation of both smooth muscle |
|----------|--|
| | and nonmuscle cell contractile activity via its phosphorylation. Implicated in cytokinesis, |
| | receptor capping, and cell locomotion. |
| | |
| | 1) Xia, Y. et al. c-Jun downregulation by HDAC3-dependent transcriptional repression promotes |
| | osmotic stress-induced cell apoptosis. Mol. Cell 25, 219–232 (2007). |
| | 2)Vander Heiden, M. G. et al. Evidence for an alternative glycolytic pathway in rapidly |
| | proliferating cells. Science 329, 1492–1499 (2010). |
| | 3) Fang, D. et al. Phosphorylation of beta-catenin by AKT promotes beta-catenin transcriptional |
| | activity. J. Biol. Chem. 282, 11221–11229 (2007). |
| | Aliases: 20 kDa myosin light chain antibody; Human 20kDa myosin light chain (MLC2) mRNA |
| | complete cds antibody; LC20 antibody; MGC3505 antibody; MLC 2 antibody; MLC-2C antibody; |
| | MLC2 antibody; MLY 9 antibody; MRLC1 antibody; MYL9 antibody; MYL9_HUMAN antibody; |
| | Myosin light chain 9 regulatory antibody; Myosin light polypeptide 9 regulatory antibody; myosin |
| | regulatory light chain 1 antibody; Myosin regulatory light chain 2 antibody; Myosin regulatory |
| | light chain 2 smooth muscle isoform antibody; Myosin regulatory light chain 9 antibody; Myosin |
| | regulatory light chain MRLC1 antibody; Myosin regulatory light polypeptide 9 antibody; Myosin |
| | RLC antibody; Myosin vascular smooth muscle light chain 2 antibody; MYRL2 antibody; |
| | OTTHUMP00000030857 antibody; smooth muscle isoform antibody |
| UniProt: | P24844 |

| Δn | nlica | tion | Details |
|----|-------|------|---------|
| Aμ | piica | UOH | Details |

| Application Notes: | WB:1:500-1:1000, IF:1:100-1:200, For Research Use only | |
|--------------------|---|--|
| Restrictions: | | |
| Handling | | |
| Format: | Liquid | |
| Buffer: | Supplied at 1.0 mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol. | |
| Preservative: | Sodium azide | |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | |
| Storage: | -20 °C,-80 °C | |
| | | |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN7138433 | 01/16/2024 | Copyright antibodies-online. All rights reserved. Storage Comment:


Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images

DAPI


MLC2 pS15





Immunofluorescence

Image 1. Immunofluorescence staining of methanol-fixed U87 cells using MLC2 (Phospho-Ser15) Antibody.



U87 cells were synchronized in I (interphase) and C (Cytokinesis) respectively, then were harvested for immunoblotting. Western Blotting

Image 2.