

Datasheet for ABIN7138753
anti-MAP2K7 antibody (pSer271)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	MAP2K7
Binding Specificity:	pSer271
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAP2K7 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Peptide sequence around phosphorylation site of Serine271(V-D-S(p)-K-A) derived from Human MAP2K7.
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using

Target Details

Target:	MAP2K7
Alternative Name:	MAP2K7 (MAP2K7 Products)

Target Details

Background:

Background:

The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase specifically activates MAPK8/JNK1 and MAPK9/JNK2, and this kinase itself is phosphorylated and activated by MAP kinase kinase kinases including MAP3K1/MEKK1, MAP3K2/MEKK2, MAP3K3/MEKK5, and MAP4K2/GCK. This kinase is involved in the signal transduction mediating the cell responses to proinflammatory cytokines, and environmental stresses. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found, but only one transcript variant has been supported and defined.

Wu Z., Mol. Cell. Biol. 17:7407-7416(1997).

Lu X., J. Biol. Chem. 272:24751-24754(1997).

Foltz I.N., J. Biol. Chem. 273:9344-9351(1998).

Aliases: c-Jun N-terminal kinase kinase 2 antibody, Dual specificity mitogen activated protein kinase kinase 7 antibody, Dual specificity mitogen-activated protein kinase kinase 7 antibody, JNK activating kinase 2 antibody, JNK kinase 2 antibody, JNK-activating kinase 2 antibody, JNKK 2 antibody, Jnkk-2 antibody, Jnkk2 antibody, MAP kinase kinase 7 antibody, MAP2K7 antibody, MAPK/ERK kinase 7 antibody, MAPKK 7 antibody, MAPKK-7 antibody, MAPKK7 antibody, MEK 7 antibody, Mitogen Activated Protein Kinase kinase 7 antibody, MKK 7 antibody, MKK-7 antibody, MKK7 antibody, MP2K7_HUMAN antibody, PRKMK 7 antibody, PRKMK-7 antibody, PRKMK7 antibody, SAPK kinase 4 antibody, SAPKK-4 antibody, SAPKK4 antibody, Sek 2 antibody, Sek-2 antibody, Sek2 antibody, SKK4 antibody, stress-activated protein kinase kinase 4 antibody

UniProt:

[O14733](#)

Pathways:

[MAPK Signaling](#), [TLR Signaling](#), [Fc-epsilon Receptor Signaling Pathway](#), [Activation of Innate immune Response](#), [Toll-Like Receptors Cascades](#), [BCR Signaling](#)

Application Details

Application Notes:

WB:1:500-1:1000, IHC:1:50-1:100,

Restrictions:

For Research Use only

Handling

Format:

Liquid


Buffer:

Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl,

Handling


	0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of paraffin-embedded human brain tissue, using MAP2K7 (Phospho-Ser271) antibody (left) or the same antibody preincubated with blocking peptide (right).



Western Blotting

Image 2. Western blot analysis of extracts from 3T3 cells treated with insulin using MAP2K7 (Phospho-Ser271) Antibody. The lane on the right is treated with the antigen-specific peptide.