

Datasheet for ABIN7138820 anti-Prohibitin antibody (pThr258)

1 Image



Go to Product page

_						
	1//	Д	rv	16	٦/	٨
U	W	\vdash	ΙV	Ιt	٦,	/V


Quantity:	100 μL
Target:	Prohibitin (PHB)
Binding Specificity:	pThr258
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Prohibitin antibody is un-conjugated
Application:	Western Blotting (WB), ELISA
Product Details	
Immunogen:	Peptide sequence around phosphorylation site of threonine 258 (N-I-T(p)-Y-L) derived from
	Human PHB.
Isotype:	IgG
Isotype: Cross-Reactivity:	IgG Human, Mouse, Rat
Cross-Reactivity:	Human, Mouse, Rat
Cross-Reactivity:	Human, Mouse, Rat Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH
Cross-Reactivity:	Human, Mouse, Rat Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific
Cross-Reactivity: Purification:	Human, Mouse, Rat Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific
Cross-Reactivity: Purification: Target Details	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi

Target Details

Background:	Background:	
	Prohibitin inhibits DNA synthesis. It has a role in regulating proliferation. As yet it is unclear if	
	the protein or the mRNA exhibits this effect. May play a role in regulating mitochondrial	
	respiration activity and in aging.	
	Aliases: Epididymis luminal protein 215 antibody; Epididymis secretory sperm binding protein Li	
	54e antibody; HEL 215 antibody; HEL S 54e antibody; PHB antibody; PHB_HUMAN antibody;	
	PHB1 antibody; Prohibitin antibody	
UniProt:	P35232	
Pathways:	Intracellular Steroid Hormone Receptor Signaling Pathway, Regulation of Intracellular Steroid	
	Hormone Receptor Signaling	

Application Details

Application Notes:	WB:1:500-1:1000,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Supplied at 1.0 mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Western Blotting

Image 1. Western blot analysis of extracts from MCF, PC12, 293 and 3T3 cells using PHB (Phospho-Thr258) Antibody.