

Datasheet for ABIN7139190  
**anti-HIST1H1C antibody (acLys62)**

## 3 Images

[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	HIST1H1C
Binding Specificity:	acLys62
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HIST1H1C antibody is un-conjugated
Application:	ELISA, Immunofluorescence (IF), Chromatin Immunoprecipitation (ChIP), Immunocytochemistry (ICC)

## Product Details

Immunogen:	Peptide sequence around site of Acetyl-Lys (62) derived from Human Histone H1.2
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Antigen Affinity Purified

## Target Details

Target:	HIST1H1C
Alternative Name:	HIST1H1C ( <a href="#">HIST1H1C Products</a> )
Background:	Background: Histone H1 protein binds to linker DNA between nucleosomes forming the

## Target Details

macromolecular structure known as the chromatin fiber. Histones H1 are necessary for the condensation of nucleosome chains into higher-order structured fibers. Acts also as a regulator of individual gene transcription through chromatin remodeling, nucleosome spacing and DNA methylation (By similarity).

Aliases: H1 histone family member 2 antibody, H1.a antibody, H12\_HUMAN antibody, H1F2 antibody, H1s-1 antibody, HIST1H1C antibody, Histone 1 H1c antibody, Histone cluster 1 H1c antibody, Histone H1.2 antibody, Histone H1c antibody, Histone H1d antibody, Histone H1s-1 antibody, MGC3992 antibody

UniProt: [P16403](#)

## Application Details

Application Notes: Recommended dilution: ICC:1:20-1:200, IF:1:50-1:200,

Restrictions: For Research Use only

## Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.




### Immunohistochemistry

**Image 1.** Chromatin Immunoprecipitation Hela ( $4 \times 10^6$ , treated with 30 mM sodium butyrate for 4h) were treated with Micrococcal Nuclease, sonicated, and immunoprecipitated with 5  $\mu$ g anti-HIST1H1C (ABIN7139190) or a control normal rabbit IgG. The resulting ChIP DNA was quantified using real-time PCR with primers against the  $\beta$ -Globin promoter.



### Immunocytochemistry

**Image 2.** Immunocytochemistry analysis of Hela cells using ABIN7139190 at dilution of 1:100



### Immunofluorescence

**Image 3.** Immunofluorescent analysis of Hela cells (sodium butyrate, 30 mM, 4h) using ABIN7139190 at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)