

Datasheet for ABIN7139191  
**anti-HIST1H1E antibody (acLys63)**

## 3 Images

[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	HIST1H1E
Binding Specificity:	acLys63
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HIST1H1E antibody is un-conjugated
Application:	ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC), Chromatin Immunoprecipitation (ChIP)

## Product Details

Immunogen:	Peptide sequence around site of Acetyl-Lys (63) derived from Human Histone H1.4
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Antigen Affinity Purified

## Target Details

Target:	HIST1H1E
Alternative Name:	HIST1H1E ( <a href="#">HIST1H1E Products</a> )
Background:	Background: Histone H1 protein binds to linker DNA between nucleosomes forming the

## Target Details

macromolecular structure known as the chromatin fiber. Histones H1 are necessary for the condensation of nucleosome chains into higher-order structured fibers. Acts also as a regulator of individual gene transcription through chromatin remodeling, nucleosome spacing and DNA methylation (By similarity).

Aliases: H1 histone family member 4 antibody, H1.4 antibody, H14\_HUMAN antibody, H1E antibody, H1F4 antibody, Hist1h1e antibody, Histone 1 H1e antibody, Histone cluster 1 H1e antibody, Histone H1 antibody, Histone H1.4 antibody, Histone H1B antibody, MGC116819 antibody

UniProt: [P10412](#)

## Application Details

Application Notes: Recommended dilution: ICC:1:20-1:200, IF:1:50-1:200,

Restrictions: For Research Use only

## Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.




### Immunohistochemistry

**Image 1.** Chromatin Immunoprecipitation Hela ( $4 \times 10^6$ , treated with 30 mM sodium butyrate for 4h) were treated with Micrococcal Nuclease, sonicated, and immunoprecipitated with 5  $\mu$ g anti-HIST1H1E (ABIN7139191) or a control normal rabbit IgG. The resulting ChIP DNA was quantified using real-time PCR with primers against the  $\beta$ -Globin promoter.



### Immunofluorescence

**Image 2.** Immunofluorescent analysis of Hela cells (sodium butyrate, 30 mM, 4h) using ABIN7139191 at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



### Immunocytochemistry

**Image 3.** Immunocytochemistry analysis of Hela cells using ABIN7139191 at dilution of 1:100