

Datasheet for ABIN7139603
anti-HIST1H1C antibody (meLys45)

3 Images

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	HIST1H1C
Binding Specificity:	meLys45
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HIST1H1C antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	Peptide sequence around site of Mono-methyl-Lys (45) derived from Human Histone H1.2
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Antigen Affinity Purified

Target Details

Target:	HIST1H1C
Alternative Name:	HIST1H1C (HIST1H1C Products)
Background:	Background: Histone H1 protein binds to linker DNA between nucleosomes forming the macromolecular structure known as the chromatin fiber. Histones H1 are necessary for the

Target Details

condensation of nucleosome chains into higher-order structured fibers. Acts also as a regulator of individual gene transcription through chromatin remodeling, nucleosome spacing and DNA methylation (By similarity).

Aliases: H1 histone family member 2 antibody, H1.a antibody, H12_HUMAN antibody, H1F2 antibody, H1s-1 antibody, HIST1H1C antibody, Histone 1 H1c antibody, Histone cluster 1 H1c antibody, Histone H1.2 antibody, Histone H1c antibody, Histone H1d antibody, Histone H1s-1 antibody, MGC3992 antibody

UniProt: [P16403](#)

Application Details

Application Notes: Recommended dilution: WB:1:100-1:1000, ICC:1:20-1:200, IF:1:10-1:100,

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C, -80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.




Immunocytochemistry

Image 1. Immunocytochemistry analysis of ABIN7139603 diluted at 1:25 and staining in HeLa cells performed on a Leica Bond™ system. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Western Blotting

Image 2. Western Blot Positive WB detected in: 293 whole cell lysate, A549 whole cell lysate. All lanes: HIST1H1C antibody at 1:100. Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 22 kDa. Observed band size: 22 kDa.



Immunofluorescence

Image 3. Immunofluorescence staining of HeLa cells with ABIN7139603 at 1:12.5, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).