

Datasheet for ABIN7139735
anti-AXIN2 antibody (AA 56-73)[Go to Product page](#)

3 Images

Overview

Quantity:	100 µL
Target:	AXIN2
Binding Specificity:	AA 56-73
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This AXIN2 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), ELISA, Immunofluorescence (IF)

Product Details

Immunogen:	Peptide sequence from Human Axin-2 protein (56-73AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Antigen Affinity Purified

Target Details

Target:	AXIN2
Alternative Name:	AXIN2 (AXIN2 Products)
Background:	Background: Inhibitor of the Wnt signaling pathway. Down-regulates beta-catenin. Probably facilitate the phosphorylation of beta-catenin and APC by GSK3B (By similarity).

Target Details

Aliases: Axil antibody, Axin like protein antibody, Axin-2 antibody, Axin-like protein antibody, Axin2 antibody, AXIN2_HUMAN antibody, Axis inhibition protein 2 antibody, Conductin antibody, DKFZp781B0869 antibody, MGC10366 antibody, MGC126582 antibody

UniProt: [Q9Y2T1](#)

Pathways: [Regulation of G-Protein Coupled Receptor Protein Signaling](#)

Application Details

Application Notes: Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200,

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images




Immunofluorescence

Image 1. Immunofluorescence staining of U251 cells with ABIN7139735 at 1:100, counter-stained with DAPI. The cells were fixed in 4 % formaldehyde, permeabilized using 0.2 % Triton X-100 and blocked in 10 % normal Goat Serum. The cells were then incubated with the antibody overnight at 4 °C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemistry

Image 2. IHC image of ABIN7139735 diluted at 1:100 and staining in paraffin-embedded human colon cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunohistochemistry

Image 3. IHC image of ABIN7139735 diluted at 1:100 and staining in paraffin-embedded human brain tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.