
antibodies - online.com







anti-DNER antibody (AA 150-168)



Image



Goto	Produ	ICT	nage

()	11/0	r\ /1	$\triangle 1 $
	$\lor \lor \vdash$	$I \vee I$	ew

Alternative Name:

Background:

Quantity:	100 μL
Target:	DNER
Binding Specificity:	AA 150-168
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DNER antibody is un-conjugated
Application:	ELISA, Immunofluorescence (IF)
Product Details	
Immunogen:	Peptide sequence from Human Delta and Notch-like epidermal growth factor-related receptor protein (150-168AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Antigen Affinity Purified
Target Details	
Target:	DNER

Background: Activator of the NOTCH1 pathway. May mediate neuron-glia interaction during

DNER (DNER Products)

astrocytogenesis (By similarity).

Aliases: bet antibody, Brain EGF repeat-containing transmembrane protein antibody, Bret antibody, Delta and Notch-like epidermal growth factor-related receptor antibody, Delta notch like EGF repeat containing transmembrane antibody, Delta/notch like EGF repeat containing antibody, Delta/notch-like EGF-related receptor antibody, Dner antibody, DNER_HUMAN antibody, PRO299 antibody, Transmembrane protein Bet antibody, UNQ26 antibody

UniProt: Q8NFT8

Pathways: Skeletal Muscle Fiber Development

Application Details

Application Notes: Recommended dilution: IF:1:50-1:200,


Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunofluorescence

Image 1. Immunofluorescence staining of U87 cells with ABIN7139803 at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).