antibodies -online.com





anti-TACC2 antibody (AA 2851-2948) (Cy7)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL	
Target:	TACC2	
Binding Specificity:	AA 2851-2948	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TACC2 antibody is conjugated to Cy7	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human TACC2
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	TACC2
Alternative Name:	TACC2 (TACC2 Products)

Target Details

rarget Details		
Background:	Synonyms: TACC2_HUMAN, Anti zuai 1, AZU 1, AZU1, ECTACC, TACC 2, Transforming acidic coiled coil containing protein 2.	
	Background: TACC2 is a conserved family of centrosome and microtubule interacting proteins	
	that are implicated in cancer. This gene encodes a protein that concentrates at centrosomes	
	throughout the cell cycle. This gene lies within a chromosomal region associated with	
	tumorigenesis. Expression of this gene is thought to affect the progression of breast tumors.	
	Expression of this gene is also induced by erythropoietin.	
Gene ID:	566	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	