antibodies -online.com





anti-TACC2 antibody (AA 2851-2948) (FITC)



()	1 /	\sim	KI /	110	Νę
	1//	\vdash	I \/	1 ←	٠// ٢

Quantity:	100 μL	
Target:	TACC2	
Binding Specificity:	AA 2851-2948	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TACC2 antibody is conjugated to FITC	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human TACC2
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	TACC2
Alternative Name:	TACC2 (TACC2 Products)

Target Details

rarget Details			
Background:	Synonyms: TACC2_HUMAN, Anti zuai 1, AZU 1, AZU1, ECTACC, TACC 2, Transforming acidic coiled coil containing protein 2.		
	Background: TACC2 is a conserved family of centrosome and microtubule interacting proteins		
	that are implicated in cancer. This gene encodes a protein that concentrates at centrosomes		
	throughout the cell cycle. This gene lies within a chromosomal region associated with		
	tumorigenesis. Expression of this gene is thought to affect the progression of breast tumors.		
	Expression of this gene is also induced by erythropoietin.		
Gene ID:	566		
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		
Expiry Date:	12 months		