# antibodies - online.com







## anti-IgG3 antibody (AA 312-324)



Image



#### Overview

Quantity:	100 μL
Target:	lgG3
Binding Specificity:	AA 312-324
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IgG3 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB)

#### **Product Details**

Immunogen:	Peptide sequence from Human Immunoglobulin heavy constant gamma 3 protein (312-324AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	lgG3
Alternative Name:	IGHG3 (IgG3 Products)
Target Type:	Antibody
Background:	Background: Constant region of immunoglobulin heavy chains. Immunoglobulins, also known

as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulins-secreting plasma cells. Secreted immunoglobulins mediate the effector phase of humoral immunity, which results in the elimination of bound antigens (PubMed:22158414, PubMed:20176268). The antigen binding site is formed by the variable domain of one heavy chain, together with that of its associated light chain. Thus, each immunoglobulin has two antigen binding sites with remarkable affinity for a particular antigen. The variable domains are assembled by a process called V-(D)-J rearrangement and can then be subjected to somatic hypermutations which, after exposure to antigen and selection, allow affinity maturation for a particular antigen (PubMed:17576170, PubMed:20176268).

Aliases: IGHG3Immunoglobulin heavy constant gamma 3 antibody, HDC antibody, Heavy chain disease protein antibody, Ig gamma-3 chain C region antibody

UniProt:


P01860

#### **Application Details**

Application Notes:

1-1	,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Recommended dilution: WB:1:1000-1:5000,



#### **Western Blotting**

**Image 1.** Western Blot Positive WB detected in human serum All lanes:IGHG3 antibody at 1:2000 Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 42 kDa Observed band size: 42 kDa