antibodies -online.com





anti-IFNA5 antibody (AA 52-73)





Go to Product page

Overview

Quantity:	100 μL
Target:	IFNA5
Binding Specificity:	AA 52-73
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IFNA5 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Peptide sequence from Human Interferon alpha-5 protein (52-73AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Antigen Affinity Purified

Target Details

Target:	IFNA5
Alternative Name:	IFNA5 (IFNA5 Products)
Background:	Background: Produced by macrophages, IFN-alpha have antiviral activities. Interferon
	stimulates the production of two enzymes: a protein kinase and an oligoadenylate synthetase.

Aliases: IeIF G antibody, IFN alphaG antibody, IFN-alpha-5 antibody, IFN-AlphaG antibody, Ifna5 antibody, IFNA5_HUMAN antibody, INA5 antibody, INFA5 antibody, Interferon alpha 61 antibody, Interferon alpha G antibody, Interferon alpha-5 antibody, Interferon alpha-61 antibody, Interferon alpha-G antibody, interferon, alpha 5 antibody, LeIF G antibody

UniProt: P01569

Pathways: JAK-STAT Signaling, Hepatitis C


Application Details

Application Notes:	Recommended dilution: IHC:1:20-1:200,
Restrictions:	For Research Use only

Handling

папишту	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7139891 diluted at 1:100 and staining in paraffin-embedded human gastric cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30 min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and



visualized using an HRP conjugated SP system.