



[Go to Product page](#)

Datasheet for ABIN7139929
anti-MDFIC antibody (AA 35-53) (Biotin)

Overview

Quantity:	100 µL
Target:	MDFIC
Binding Specificity:	AA 35-53
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MDFIC antibody is conjugated to Biotin
Application:	ELISA

Product Details

Immunogen:	Peptide sequence from Human MyoD family inhibitor domain-containing protein (35-53AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	MDFIC
Alternative Name:	MDFIC (MDFIC Products)
Background:	Background: Acts as a transcriptional activator or repressor. Inhibits the transcriptional activation of Zic family proteins ZIC1, ZIC2 and ZIC3. Retains nuclear Zic proteins ZIC1, ZIC2

Target Details

and ZIC3 in the cytoplasm. Modulates the expression from both cellular and viral promoters. Down-regulates Tat-dependent transcription of the human immunodeficiency virus type 1 (HIV-1) LTR by interacting with HIV-1 Tat and Rev and impairing their nuclear import, probably by rendering the NLS domains inaccessible to importin-beta. Also stimulates activation of human T-cell leukemia virus type I (HTLV-I) LTR. Binds to the axin complex, resulting in an increase in the level of free beta-catenin. Affects axin regulation of the WNT and JNK signaling pathways. Aliases: MDFIC antibody, MyoD family inhibitor domain-containing protein antibody, I-mfa domain-containing protein antibody, hIC antibody

UniProt: [Q9P1T7](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C, -80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.