antibodies .- online.com







anti-NPFFR2 antibody (AA 23-43) (FITC)



Overview

Quantity:	100 μL
Target:	NPFFR2 (NPFF2)
Binding Specificity:	AA 23-43
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NPFFR2 antibody is conjugated to FITC
Application:	Please inquire

Product Details

Immunogen:	Peptide sequence from Human Neuropeptide FF receptor 2 protein (23-43AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	NPFFR2 (NPFF2)
Alternative Name:	NPFFR2 (NPFF2 Products)
Background:	Background: Receptor for NPAF (A-18-F-amide) and NPFF (F-8-F-amide) neuropeptides, also known as morphine-modulating peptides. Can also be activated by a variety of naturally

occurring or synthetic FMRF-amide like ligands. This receptor mediates its action by association with G proteins that activate a phosphatidylinositol-calcium second messenger system.

Aliases: G protein coupled receptor 74 antibody, G protein coupled receptor HLWAR 77 antibody, G protein coupled receptor HLWAR77 antibody, G-protein coupled receptor 74 antibody, G-protein coupled receptor HLWAR77 antibody, GPR 74 antibody, GPR74 antibody, HLWAR 77 antibody, Neuropeptide FF 2 antibody, Neuropeptide FF receptor 2 antibody, Neuropeptide G protein coupled receptor antibody, Neuropeptide G-protein coupled receptor antibody, NPFF 2 antibody, NPFF 2 antibody, NPFF 2 antibody, NPFF2 antibody, NPFF2 antibody, NPFF2 antibody, NPFF2 antibody

UniProt:

Q9Y5X5

Pathways:

cAMP Metabolic Process

Application Details

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.