antibodies .- online.com







anti-GRF2 antibody (AA 22-40) (HRP)



Go to Product page

| () | 1/0 | r\ /1 | 014 | |
|-----|-----|-------|-----|---|
| () | ve | I V I | -v | V |
| | | | | |

| Quantity: | 100 μL |
|----------------------|---|
| Target: | GRF2 (RAPGEF1) |
| Binding Specificity: | AA 22-40 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This GRF2 antibody is conjugated to HRP |
| Application: | ELISA |

Product Details

| Immunogen: | Peptide sequence from Human Rap guanine nucleotide exchange factor 1 protein (22-40AA) |
|-------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | GRF2 (RAPGEF1) |
|-------------------|---|
| Alternative Name: | RAPGEF1 (RAPGEF1 Products) |
| Background: | Background: Guanine nucleotide-releasing protein that binds to SH3 domain of CRK and |
| | GRB2/ASH. Transduces signals from CRK to activate RAS. Plays a role in the establishment of |

| basal endoth | elial barrier function | . Plays a role in | nerve growth | factor (NGF)-ir | nduced sus | tained |
|---------------|------------------------|-------------------|--------------|-----------------|------------|--------|
| activation of | Rap1 and neurite ou | tgrowth. | | | | |

Aliases: C3G antibody, C3G protein antibody, CRK SH3 binding GNRP antibody, CRK SH3-binding GNRP antibody, DKFZp781P1719 antibody, GRF 2 antibody, GRF2 antibody, Guanine nucleotide releasing factor 2 (specific for crk proto oncogene) antibody, Guanine nucleotide releasing factor 2 antibody, Guanine nucleotide-releasing factor 2 antibody, OTTHUMP00000064558 antibody, Protein C3G antibody, Rap guanine nucleotide exchange

OTTHUMP00000064558 antibody, Protein C3G antibody, Rap guanine nucleotide exchange factor (GEF) 1 antibody, Rap guanine nucleotide exchange factor 1 antibody, RAPGEF 1 antibody, RAPGEF1 antibody, RAPGEF1 antibody

UniProt: Q13905

Pathways: Interferon-gamma Pathway, Neurotrophin Signaling Pathway, Platelet-derived growth Factor

Receptor Signaling, Signaling of Hepatocyte Growth Factor Receptor

Application Details

| Application Notes: | Optimal working dilution should be determined by the investigator. |
|--------------------|--|
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid | |
|--------------------|---|--|
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4 | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | |
| Storage: | -20 °C,-80 °C | |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. | |