



[Go to Product page](#)

Datasheet for ABIN7140124  
**anti-LEO1 antibody (AA 2-24) (FITC)**

### Overview

Quantity:	100 µL
Target:	LEO1
Binding Specificity:	AA 2-24
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LEO1 antibody is conjugated to FITC
Application:	Please inquire

### Product Details

Immunogen:	Peptide sequence from Human RNA polymerase-associated protein LEO1 protein (2-24AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	LEO1
Alternative Name:	LEO1 ( <a href="#">LEO1 Products</a> )
Background:	Background: Component of the PAF1 complex (PAF1C) which has multiple functions during transcription by RNA polymerase II and is implicated in regulation of development and

## Target Details

---

maintenance of embryonic stem cell pluripotency. PAF1C associates with RNA polymerase II through interaction with POLR2A CTD non-phosphorylated and 'Ser-2'- and 'Ser-5'-phosphorylated forms and is involved in transcriptional elongation, acting both independently and synergistically with TCEA1 and in cooperation with the DSIF complex and HTATSF1. PAF1C is required for transcription of Hox and Wnt target genes. PAF1C is involved in hematopoiesis and stimulates transcriptional activity of KMT2A/MLL1, it promotes leukemogenesis through association with KMT2A/MLL1-rearranged oncoproteins, such as KMT2A/MLL1-MLLT3/AF9 and KMT2A/MLL1-MLLT1/ENL. PAF1C is involved in histone modifications such as ubiquitination of histone H2B and methylation on histone H3 'Lys-4' (H3K4me3). PAF1C recruits the RNF20/40 E3 ubiquitin-protein ligase complex and the E2 enzyme UBE2A or UBE2B to chromatin which mediate monoubiquitination of 'Lys-120' of histone H2B (H2BK120ub1), UB2A/B-mediated H2B ubiquitination is proposed to be coupled to transcription. PAF1C is involved in mRNA 3' end formation probably through association with cleavage and poly(A) factors. In case of infection by influenza A strain H3N2, PAF1C associates with viral NS1 protein, thereby regulating gene transcription. Involved in polyadenylation of mRNA precursors. Connects PAF1C to Wnt signaling.

Aliases: Gm185 antibody, LEO1 antibody, LEO1\_HUMAN antibody, LOC123439 antibody, Paf1/RNA polymerase II complex component antibody, Replicative senescence down regulated leo1 like protein antibody, Replicative senescence down-regulated leo1-like protein antibody, Replicative senescence downregulated leo1 like antibody, RNA polymerase associated protein LEO1 antibody, RNA polymerase-associated protein LEO1 antibody, Senescence downregulated leo1 like antibody

---

UniProt: [Q8WVC0](#)

---

Pathways: [Stem Cell Maintenance](#)

## Application Details

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

---

Preservative: ProClin

---

## Handling

---

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.