# antibodies -online.com





### Datasheet for ABIN7140246

# anti-GLI2 antibody (AA 2-20) (HRP)



#### Overview

Quantity:	100 μL
Target:	GLI2
Binding Specificity:	AA 2-20
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GLI2 antibody is conjugated to HRP
Application:	ELISA

#### **Product Details**

Immunogen:	Peptide sequence from Human Zinc finger protein GLI2 protein (2-20AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Antigen Affinity Purified

## Target Details

Target:	GLI2
Alternative Name:	GLI2 (GLI2 Products)
Background:	Background: Functions as transcription regulator in the hedgehog (Hh) pathway
	(PubMed:18455992). Functions as transcriptional activator (PubMed:9557682,

PubMed:19878745, PubMed:24311597). May also function as transcriptional repressor (By similarity). Requires STK36 for full transcriptional activator activity. Required for normal embryonic development (PubMed:15994174, PubMed:20685856).

Aliases: CJS antibody, Gli 2 antibody, GLI family zinc finger 2 antibody, GLI Kruppel family member GLI2 antibody, GLI2 antibody, GLI2\_HUMAN antibody, Glioma associated oncogene family zinc finger antibody, HPE9 antibody, Oncogene GLI2 antibody, PHS2 antibody, Tax helper protein 1 antibody, Tax helper protein 2 antibody, Tax helper protein antibody, Tax responsive element 2 holding protein antibody, Tax responsive element 25 bp sequence binding protein antibody, THP1 antibody, THP2 antibody, Zinc finger protein GLI2 antibody

UniProt: P10070

Pathways: Hedgehog Signaling, Dopaminergic Neurogenesis

#### **Application Details**

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

#### Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.