antibodies .- online.com





anti-ADCY1 antibody (AA 1-358) (FITC)



Overview

Quantity:	100 μg
Target:	ADCY1
Binding Specificity:	AA 1-358
Reactivity:	African cassava mosaic virus (ACMV)
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ADCY1 antibody is conjugated to FITC
Application:	Please inquire

Product Details

Immunogen:	Recombinant African cassava mosaic virus Replication-associated protein (1-358AA)
Isotype:	IgG
Cross-Reactivity:	African cassava mosaic virus (ACMV)
Purification:	>95%, Protein G purified

Target Details

Target:	ADCY1
Alternative Name:	AC1 (ADCY1 Products)
Background:	Background: Essential for the replication of viral ssDNA. The closed circular ssDNA genome is
	first converted to a superhelical dsDNA. Rep binds a specific region at the genome origin of

replication. It introduces an endonucleolytic nick within the conserved sequence 5\'TAATATTAC-3\' in the intergenic region of the genome present in all geminiviruses, thereby initiating the rolling circle replication (RCR). Following cleavage, binds covalently to the 5\'phosphate of DNA as a tyrosyl ester. The cleavage gives rise to a free 3\'-OH that serves as a primer for the cellular DNA polymerase. The polymerase synthesizes the (+) strand DNA by rolling circle mechanism. After one round of replication, a Rep-catalyzed nucleotidyl transfer reaction releases a circular single-stranded virus genome, thereby terminating the replication. Displays origin-specific DNA cleavage, nucleotidyl transferase, ATPase and helicase activities (By similarity).

Aliases: AC1 antibody, AL1 antibody, Replication-associated protein antibody, Rep antibody, EC 2.7.7.- antibody, EC 3.1.21.- antibody, 40.4 kDa protein antibody, Protein AC1 antibody, Protein AL1 antibody

UniProt:

P14982

Pathways:

EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Thyroid Hormone Synthesis, cAMP Metabolic Process, Myometrial Relaxation and Contraction, G-protein mediated Events, Interaction of EGFR with phospholipase C-gamma

Application Details

Restrictions:

For Research Use only

Handling

Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	