

# Datasheet for ABIN7140810 anti-EPO antibody (AA 26-192)

# 1 Image



#### Overview

| Overview             |                                    |
|----------------------|------------------------------------|
| Quantity:            | 100 μg                             |
| Target:              | EPO                                |
| Binding Specificity: | AA 26-192                          |
| Reactivity:          | Cow                                |
| Host:                | Rabbit                             |
| Clonality:           | Polyclonal                         |
| Conjugate:           | This EPO antibody is un-conjugated |
| Application:         | Western Blotting (WB), ELISA       |
| Product Details      |                                    |

| Immunogen:        | Recombinant Bovine Erythropoietin protein (26-192AA) |
|-------------------|--|
| Isotype:          | IgG  |
| Cross-Reactivity: | Cow, Human   |
| Purification:     | >95%, Protein G purified                             |

## Target Details

| Target:           | EPO                |
|-------------------|--------------------|
| Alternative Name: | EPO (EPO Products) |
| Target Type:      | Hormone            |
|                   |                    |


#### **Target Details**

|           | differentiation and the maintenance of a physiological level of circulating erythrocyte mass.  Aliases: EPOErythropoietin antibody      |
|-----------|---|
| UniProt:  | P48617  |
| Pathways: | JAK-STAT Signaling, Hormone Activity, Negative Regulation of intrinsic apoptotic Signaling, Negative Regulation of Transporter Activity |

### **Application Details**

| Application Notes: | Recommended dilution: WB:1:1000-1:5000,   |
|--------------------|---|
| Restrictions:      | For Research Use only   |
| Handling           |   |
| Format:            | Liquid  |
| Buffer:            | Preservative: 0.03 % Proclin 300<br>Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4                                |
| Preservative:      | ProClin   |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage:           | -20 °C,-80 °C   |
| Storage Comment:   | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.   |

#### **Images**



#### **Western Blotting**

**Image 1.** Western blot All lanes: EPO antibody at  $3 \mu g/mL + HepG2$  whole cell lysate Secondary Goat polyclonal to rabbit IgG at 1/10000 dilution Predicted band size: 22 kDa Observed band size: 22 kDa