

## Datasheet for ABIN7140914

## anti-LC3B antibody (AA 1-116) (FITC)



Go to Product page

_					
	W	0	rv	10	W

Quantity:	100 μg	
Target:	LC3B (MAP1LC3B)	
Binding Specificity:	AA 1-116	
Reactivity:	Candida sp.	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This LC3B antibody is conjugated to FITC	
Application:	Please inquire	
Product Details		
Immunogen:	Recombinant Candida glabrata Autophagy-related protein 8 protein (1-116AA)	
Isotype:	IgG	
Cross-Reactivity:	Candida sp.	
Purification:	>95%, Protein G purified	
Target Details		
Target:	LC3B (MAP1LC3B)	
	LC3B (MAP1LC3B)  ATG8 (MAP1LC3B Products)	

autophagosomes to the vacuole via the microtubule cytoskeleton. Required for selective autophagic degradation of the nucleus (nucleophagy) as well as for mitophagy which contributes to regulate mitochondrial quantity and quality by eliminating the mitochondria to a basal level to fulfill cellular energy requirements and preventing excess ROS production. Participates also in membrane fusion events that take place in the early secretory pathway. Also involved in endoplasmic reticulum-specific autophagic process and is essential for the survival of cells subjected to severe ER stress. The ATG8-PE conjugate mediates tethering between adjacent membranes and stimulates membrane hemifusion, leading to expansion of the autophagosomal membrane during autophagy.

Aliases: ATG8 antibody, AUT7 antibody, CAGL0A04675gAutophagy-related protein 8 antibody,

Aliases: ATG8 antibody, AUT7 antibody, CAGL0A04675gAutophagy-related protein 8 antibody, Autophagy-related ubiquitin-like modifier ATG8 antibody

UniProt:

O6FXR8

Pathways:

Autophagy

## **Application Details**

Restrictions:

For Research Use only

## Handling

Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	