antibodies - online.com







anti-OMP antibody (AA 23-394)



Image



()	1 /	\sim	KI /	110	Νę
	1//	\vdash	I \/	1 ←	٠// ٢

Quantity:	100 μL
Target:	OMP
Binding Specificity:	AA 23-394
Reactivity:	Chlamydia
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OMP antibody is un-conjugated
Application:	ELISA, Western Blotting (WB)

Product Details

lmmunogen:	Recombinant Chlamydia trachomatis serovar L2 Major outer membrane porin protein (23-394AA)
Isotype:	IgG
Cross-Reactivity:	Chlamydia
Purification:	>95%, Protein G purified

Target Details

Target:	OMP
Alternative Name:	ompA (OMP Products)
Background:	Background: In elementary bodies (EBs, the infectious stage, which is able to survive outside

Target Details

the host cell) provides the structural integrity of the outer envelope through disulfide cross-links with the small cysteine-rich protein and the large cysteine-rich periplasmic protein. It has been described in publications as the Sarkosyl-insoluble COMC (Chlamydia outer membrane complex), and serves as the functional equivalent of peptidoglycan (By similarity). Aliases: ompA antibody, omp1L2 antibody, CTL0050 antibody, Major outer membrane porin antibody, MOMP antibody

UniProt:

Buffer:

P06597


Application Details

Application Notes:	Recommended dilution: WB:1:500-1:5000,	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	

Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: 0.03 % Proclin 300

	Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C



Western Blotting

Image 1. Western Blot Positive WB detected in Recombinant protein All lanes: ompA antibody at $3.6~\mu$ g/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 58~kDa