
# antibodies - online.com







# anti-PLCD4 antibody (AA 1-762) (Biotin)



( )	11/0	K\ /	iew	1
	$\cup$	ועוי	$\square \vee \vee$	ı

Quantity:	100 μg
Target:	PLCD4
Binding Specificity:	AA 1-762
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PLCD4 antibody is conjugated to Biotin
Application:	ELISA

#### **Product Details**

lmmunogen:	Recombinant Human 1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase delta-4 protein (1-762AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## **Target Details**

Target:	PLCD4
Alternative Name:	PLCD4 (PLCD4 Products)
Background: Background: Hydrolyzes the phosphatidylinositol 4,5-bisphosphate (PIP2) to generate 2 se	

messenger molecules diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3). DAG mediates the activation of protein kinase C (PKC), while IP3 releases Ca2+ from intracellular stores. Required for acrosome reaction in sperm during fertilization, probably by acting as an important enzyme for intracellular Ca2+ mobilization in the zona pellucida-induced acrosome reaction. May play a role in cell growth. Modulates the liver regeneration in cooperation with nuclear PKC. Overexpression up-regulates the Erk signaling pathway and proliferation.

Aliases: 1 phosphatidylinositol 4 5 bisphosphate phosphodiesterase delta 4 antibody, 1-phosphatidylinositol 4 antibody, 1-phosphatidylinositol-4 antibody, 5-bisphosphate phosphodiesterase delta-4 antibody, hPLCD4 antibody, MGC12837 antibody, Phosphoinositide phospholipase C delta 4 antibody, Phospholipase C-delta-4 antibody, PLC delta 4 antibody

UniProt: Q9BRC7

Pathways: Inositol Metabolic Process

### **Application Details**

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

#### Handling

Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	