antibodies .- online.com





anti-PSMD2 antibody (Regulatory Subunit 2) (FITC)



Go to Product page

()	11/0	K\ /	iew
	\cup	ועוי	$I \cap VV$

Quantity:	100 μg
Target:	PSMD2
Binding Specificity:	AA 401-700, Regulatory Subunit 2
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PSMD2 antibody is conjugated to FITC
Application:	Please inquire

Product Details

Immunogen:	Recombinant Human 26S proteasome non-ATPase regulatory subunit 2 protein (401-700AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	

Target Details

Target:	PSMD2
Alternative Name:	PSMD2 (PSMD2 Products)
Background:	Background: Acts as a regulatory subunit of the 26 proteasome which is involved in the ATP-dependent degradation of ubiquitinated proteins.

Aliases: 26S proteasome non-ATPase regulatory subunit 2 antibody, 26S proteasome regulatory subunit RPN1 antibody, 26S proteasome regulatory subunit S2 antibody, 26S proteasome subunit p97 antibody, 55.11 protein antibody, MGC14274 antibody, P97 antibody, Proteasome (prosome macropain) 26S subunit non ATPase 2 antibody, Proteasome 26S subunit, non-ATPase, 2 antibody, Proteasome 26S, subunit 2 antibody, Protein 55.11 antibody, Psmd2 antibody, PSMD2_HUMAN antibody, Rpn1 antibody, S2 antibody, TNFR associated protein 2 antibody, TRAP2 antibody, Tumor necrosis factor receptor associated protein 2 antibody, Tumor necrosis factor type 1 receptor-associated protein 2 antibody

UniProt:

Q13200

Pathways:

Mitotic G1-G1/S Phases, DNA Replication, Synthesis of DNA, Ubiquitin Proteasome Pathway

Application Details

Restrictions:

For Research Use only

Handling

Handling		
Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	