antibodies .- online.com





anti-ABI2 antibody (AA 164-452) (HRP)



Overview

Quantity:	100 μg
Target:	ABI2
Binding Specificity:	AA 164-452
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ABI2 antibody is conjugated to HRP
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Abl interactor 2 protein (164-452AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	ABI2
Alternative Name:	ABI2 (ABI2 Products)
Background:	Background: May act in regulation of cell growth and transformation by interacting with
	nonreceptor tyrosine kinases ABL1 and/or ABL2. Part of the WAVE complex that regulates

lamellipodia formation. The WAVE complex regulates actin filament reorganization via its interaction with the Arp2/3 complex. Regulates ABL1/c-Abl-mediated phosphorylation of MENA.

Aliases: Abelson interactor 2 antibody, ABI 2 antibody, Abi-2 antibody, ABI2 antibody, ABI2_HUMAN antibody, ABI2B antibody, AbI binding protein 3 antibody, AbI interacting protein 1 (SH3 containing protein) antibody, AbI interactor 2 antibody, AbI interactor protein 2b antibody, AbI-binding protein 3 antibody, AbIBP3 antibody, AIP 1 antibody, Arg binding protein 1 antibody, Arg protein tyrosine kinase binding protein antibody, Arg-binding protein 1 antibody, ArgBP1 antibody, ARGBPIA antibody, ArgBPIB antibody, OTTHUMP00000163783 antibody, OTTHUMP00000163784 antibody, OTTHUMP00000206182 antibody, OTTHUMP00000206183 antibody, OTTHUMP00000206184 antibody, OTTHUMP00000206185 antibody, SSH3BP2 antibody

UniProt:

Q9NYB9

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Buffer:
Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative:
ProClin

Precaution of Use:
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage:
-20 °C,-80 °C

Upon receipt, store at -20 °C or -80 °C. Avoid repeated freeze.