antibodies -online.com





anti-SCD antibody (AA 10-70)





Overview

| Quantity: | 100 μg |
|----------------------|------------------------------------|
| Target: | SCD |
| Binding Specificity: | AA 10-70 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This SCD antibody is un-conjugated |
| Application: | ELISA, Immunohistochemistry (IHC) |

Product Details

| Immunogen: | Recombinant Human Acyl-CoA desaturase protein (10-70AA) |
|-------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | SCD |
|-------------------|--|
| Alternative Name: | SCD (SCD Products) |
| Background: | Background: Stearyl-CoA desaturase that utilizes O(2) and electrons from reduced cytochrome b5 to introduce the first double bond into saturated fatty acyl-CoA substrates |

(PubMed:15907797, PubMed:18765284). Catalyzes the insertion of a cis double bond at the delta-9 position into fatty acyl-CoA substrates including palmitoyl-CoA and stearoyl-CoA (PubMed:15907797, PubMed:18765284). Gives rise to a mixture of 16:1 and 18:1 unsaturated fatty acids (PubMed:15610069). Plays an important role in lipid biosynthesis. Plays an important role in regulating the expression of genes that are involved in lipogenesis and in regulating mitochondrial fatty acid oxidation (By similarity). Plays an important role in body energy homeostasis (By similarity). Contributes to the biosynthesis of membrane phospholipids, cholesterol esters and triglycerides (By similarity).

Aliases: ACOD_HUMAN antibody, Acyl-CoA desaturase antibody, Delta(9)-desaturase antibody, Delta-9 desaturase antibody, Fatty acid desaturase antibody, PR00998 antibody, SCD 1 antibody, SCD antibody, SCD1 antibody,

UniProt:

000767

Pathways:

Brown Fat Cell Differentiation

Stearoyl-CoA desaturase antibody

Application Details

Application Notes:


Recommended dilution: IHC:1:20-1:200,

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|---|
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |



Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human lung cancer using ABIN7143196 at dilution of 1:100