



[Go to Product page](#)

Datasheet for ABIN7143604
anti-AGRN antibody (AA 968-1130) (HRP)

Overview

Quantity:	100 µg
Target:	AGRN
Binding Specificity:	AA 968-1130
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This AGRN antibody is conjugated to HRP
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Agrin protein (968-1130AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	AGRN
Alternative Name:	AGRN (AGRN Products)
Background:	Background: Isoform 1: heparan sulfate basal lamina glycoprotein that plays a central role in the formation and the maintenance of the neuromuscular junction (NMJ) and directs key

Target Details

events in postsynaptic differentiation. Component of the AGRN-LRP4 receptor complex that induces the phosphorylation and activation of MUSK. The activation of MUSK in myotubes induces the formation of NMJ by regulating different processes including the transcription of specific genes and the clustering of AChR in the postsynaptic membrane. Calcium ions are required for maximal AChR clustering. AGRN function in neurons is highly regulated by alternative splicing, glycan binding and proteolytic processing. Modulates calcium ion homeostasis in neurons, specifically by inducing an increase in cytoplasmic calcium ions. Functions differentially in the central nervous system (CNS) by inhibiting the alpha(3)-subtype of Na⁺/K⁺-ATPase and evoking depolarization at CNS synapses. This secreted isoform forms a bridge, after release from motor neurons, to basal lamina through binding laminin via the NtA domain.

Aliases: AGRIN antibody, Agrin C-terminal 22 kDa fragment antibody, Agrin proteoglycan antibody, AGRIN_HUMAN antibody, Agrn antibody, C22 antibody, C90 antibody, FLJ45064 antibody, OTTHUMP00000044043 antibody

UniProt: [O00468](#)

Pathways: [Glycosaminoglycan Metabolic Process](#), [Regulation of Muscle Cell Differentiation](#), [Skeletal Muscle Fiber Development](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C, -80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.