antibodies -online.com







anti-RAI14 antibody (AA 242-346)





Go to Product page

| \sim | |
|--------|-----------|
| ()\/白 | view |
| | V I C V V |

| Quantity: | 100 μg |
|----------------------|--------------------------------------|
| Target: | RAI14 |
| Binding Specificity: | AA 242-346 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This RAI14 antibody is un-conjugated |
| Application: | ELISA, Immunohistochemistry (IHC) |

Product Details

| Immunogen: | Recombinant Human Ankycorbin protein (242-346AA) |
|-------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | RAI14 |
|-------------------|---|
| Alternative Name: | RAI14 (RAI14 Products) |
| Background: | Background: cytosol, fibrillar center, nucleus |
| | Aliases: Ankycorbin antibody, Ankyrin repeat and coiled coil structure containing protein |

Target Details

antibody, Ankyrin repeat and coiled-coil structure-containing protein antibody, DKFZp564G013 antibody, KIAA1334 antibody, NORPEG antibody, Novel retinal pigment epithelial cell protein antibody, RAI13 antibody, RAI14 antibody, RAI14_HUMAN antibody, Retinoic acid induced 14 antibody, Retinoic acid induced protein 14 antibody

UniProt:

Q9P0K7


Application Details

| Application Notes: | Recommended dilution: IHC:1:500-1:1000, |
|--------------------|---|
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid |
|--------------------|---|
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |

Images



Immunohistochemistry

Image 1. IHC image of ABIN7144147 diluted at 1:700 and staining in paraffin-embedded human endometrial cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30 min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.