antibodies .- online.com





anti-TAP2 antibody (AA 473-615) (FITC)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	ועוי	$\square \vee \vee$	ı

Quantity:	100 μg
Target:	TAP2
Binding Specificity:	AA 473-615
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TAP2 antibody is conjugated to FITC
Application:	Please inquire

Product Details

Immunogen:	Recombinant Human Antigen peptide transporter 2 protein (473-615AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	

Target Details

Target:	TAP2	
Alternative Name:	TAP2 (TAP2 Products)	
Background:	Background: Involved in the transport of antigens from the cytoplasm to the endoplasmic	
	reticulum for association with MHC class I molecules. Also acts as a molecular scaffold for the	

final stage of MHC class I folding, namely the binding of peptide. Nascent MHC class I molecules associate with TAP via tapasin. Inhibited by the covalent attachment of herpes simplex virus ICP47 protein, which blocks the peptide-binding site of TAP. Inhibited by human cytomegalovirus US6 glycoprotein, which binds to the lumenal side of the TAP complex and inhibits peptide translocation by specifically blocking ATP-binding to TAP1 and prevents the conformational rearrangement of TAP induced by peptide binding. Inhibited by human adenovirus E3-19K glycoprotein, which binds the TAP complex and acts as a tapasin inhibitor, preventing MHC class I/TAP association.

Aliases: ABC transporter, MHC 2 antibody, ABC18 antibody, ABCB3 antibody, Antigen peptide transporter 2 antibody, APT2 antibody, ATP binding cassette, sub family B (MDR/TAP), member 3 antibody, ATP-binding cassette sub-family B member 3 antibody, D6S217E antibody, Peptide supply factor 2 antibody, Peptide transporter involved in antigen processing 2 antibody, Peptide transporter PSF2 antibody, Peptide transporter TAP2 antibody, PSF 2 antibody, PSF-2 antibody, PSF-2 antibody, Really interesting new gene 11 protein antibody, RING 11 antibody, RING11 antibody, TAP 2 antibody, Tap2 antibody, TAP2_HUMAN antibody, Transporter 2 ATP binding cassette sub family B antibody, Transporter 2, ABC (ATP binding cassette antibody, Transporter 2, ATP binding cassette, sub family B (MDR/TAP) antibody

UniProt:

Q03519

Pathways:

Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Human Leukocyte Antigen (HLA) in Adaptive Immune Response

Application Details

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C

Storage Comment:

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.