# antibodies .- online.com





### Datasheet for ABIN714452

# anti-Epigen antibody (AA 51-130) (Cy7)



#### Overview

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | Epigen (EPGN)  |
| Binding Specificity: | AA 51-130  |
| Reactivity:          | Human, Mouse   |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This Epigen antibody is conjugated to Cy7  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human Epigen |
|-----------------------|--|
| Isotype:              | IgG  |
| Cross-Reactivity:     | Human, Mouse   |
| Predicted Reactivity: | Rat,Dog,Cow,Sheep,Horse                                    |
| Purification:         | Purified by Protein A.                                     |

## **Target Details**

| Target:           | Epigen (EPGN)          |
|-------------------|------------------------|
| Alternative Name: | Epigen (EPGN Products) |

#### **Target Details**

| rarget Details      |   |
|---------------------|---|
| Background:         | Synonyms: EPG, Epigen precursor, Epithelial mitogen, EPGN_HUMAN.  Background: Epigen is an EGF-related polypeptide growth factor that signals through the ErbB receptor-1. It is produced in several tissues, including the testis, liver, heart and in certain tumor cells. Epigen is mitogenic for fibroblasts and epithelial cells. Human Epigen is initially synthesized as a glycosylated 14.7 kDa transmembrane precursor protein, which is processed by proteolytic cleavage to produce a mature soluble sequence. |
| Gene ID:            | 255324  |
| Pathways:           | RTK Signaling, EGFR Signaling Pathway   |
| Application Details |   |
| Application Notes:  | IF(IHC-P) 1:50-200<br>IF(IHC-F) 1:50-200<br>IF(ICC) 1:50-200  |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Liquid  |
| Concentration:      | 1 μg/μL   |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.  |
| Preservative:       | ProClin   |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.  |
| Storage:            | -20 °C  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.   |
| Expiry Date:        | 12 months   |