antibodies -online.com





anti-Epigen antibody (AA 51-130) (FITC)

Datasheet for ABIN714453

Go to Product page

_						
	\/		r١	/1	0	۱۸/
\sim	V	$\overline{}$	1 V	١.	\sim	VΥ

Quantity:	100 μL
Target:	Epigen (EPGN)
Binding Specificity:	AA 51-130
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Epigen antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Epigen
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Sheep,Horse
Purification:	Purified by Protein A.

Target Details

Target:	Epigen (EPGN)	
Alternative Name:	Epigen (EPGN Products)	

Target Details

. a. got 2 otamo	
Background:	Synonyms: EPG, Epigen precursor, Epithelial mitogen, EPGN_HUMAN.
	Background: Epigen is an EGF-related polypeptide growth factor that signals through the ErbB
	receptor-1. It is produced in several tissues, including the testis, liver, heart and in certain tumor
	cells. Epigen is mitogenic for fibroblasts and epithelial cells. Human Epigen is initially
	synthesized as a glycosylated 14.7 kDa transmembrane precursor protein, which is processed
	by proteolytic cleavage to produce a mature soluble sequence.
Gene ID:	255324
Pathways:	RTK Signaling, EGFR Signaling Pathway
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months