antibodies .- online.com





anti-ARID1A antibody (AA 655-816) (HRP)



Go to Product page

| \sim | | | | | |
|--------|----|----|----|-------------|---|
| () | VE | ۲۱ | /1 | \triangle | Λ |

| Quantity: | 100 μg |
|----------------------|---|
| Quantity. | 100 μg |
| Target: | ARID1A |
| Binding Specificity: | AA 655-816 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This ARID1A antibody is conjugated to HRP |
| Application: | ELISA |

Product Details

| Immunogen: | Recombinant Human AT-rich interactive domain-containing protein 1A protein (655-816AA) |
|-------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | ARID1A |
|-------------------|--|
| Alternative Name: | ARID1A (ARID1A Products) |
| Background: | Background: Involved in transcriptional activation and repression of select genes by chromatin |
| | remodeling (alteration of DNA-nucleosome topology). Binds DNA non-specifically. Belongs to |

the neural progenitors-specific chromatin remodeling complex (npBAF complex) and the neuron-specific chromatin remodeling complex (nBAF complex). During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural stem/progenitor cells to post-mitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the self-renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along with CREST plays a role regulating the activity of genes essential for dendrite growth (By similarity).

Aliases: actin-dependent regulator of chromatin subfamily F member 1 antibody, ARID domain-containing protein 1A antibody, ARID domain-containing protein 1A antibody, ARID domain-containing protein 1A antibody, ARID1A antibody, AT rich interactive domain 1A (SWI like) antibody, AT rich interactive domain 1A antibody, AT rich interactive domain containing protein 1A antibody, AT-rich interactive domain-containing protein 1A antibody, B120 antibody, BAF250 antibody, BAF250 antibody, BAF250A antibody, BM029 antibody, brain protein 120 antibody, BRG1 associated factor 250 antibody, BRG1-associated factor 250 antibody, BRG1-associated factor 250 antibody, BRG1-associated factor 250 antibody, C10RF4 antibody, chromatin remodeling factor p250 antibody, chromosome 1 open reading frame 4 antibody, ELD antibody, hELD antibody, hOSA1 antibody, matrix-associated antibody, MRD14 antibody, Osa homolog 1 antibody, OSA1 antibody, OSA1 nuclear protein antibody, P270 antibody, SMARCF1 antibody, SWI like protein antibody, SWI SNF complex protein p270 antibody, SWI-like protein antibody, SWI/SNF complex protein p270 antibody, SWI-SNF related, matrix associated, actin dependent regulator of chromatin, subfamily f, member 1 antibody, SWI/SNF-related antibody

UniProt:

014497

Pathways:

Intracellular Steroid Hormone Receptor Signaling Pathway, Tube Formation

Application Details

Application Notes:

Optimal working dilution should be determined by the investigator.

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|---|
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |