# antibodies - online.com







## anti-ATP5F1D antibody (AA 32-158)



**Images** 



$\sim$	
( )\/\Di	view
	VICVV

Quantity:	100 μg
Target:	ATP5F1D
Binding Specificity:	AA 32-158
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ATP5F1D antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (IHC)

#### **Product Details**

Immunogen:	Recombinant Human ATP synthase subunit delta, mitochondrial protein (32-158AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	ATP5F1D
Alternative Name:	ATP5F1D (ATP5F1D Products)
Background:	Background: Mitochondrial membrane ATP synthase (F1F0 ATP synthase or Complex V)
	produces ATP from ADP in the presence of a proton gradient across the membrane which is

generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F1 - containing the extramembraneous catalytic core, and F0 - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP turnover in the catalytic domain of F1 is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Part of the complex F1 domain and of the central stalk which is part of the complex rotary element. Rotation of the central stalk against the surrounding alpha3beta3 subunits leads to hydrolysis of ATP in three separate catalytic sites on the beta subunits.

Aliases: ATP synthase subunit delta, mitochondrial antibody, ATP synthase subunit delta, mitochondrial antibody, ATP synthase, H+ transporting, mitochondrial F1 complex, delta subunit antibody, ATP5D antibody, ATPD\_HUMAN antibody, F ATPase delta subunit antibody, Mitochondrial ATP synthase complex delta subunit precusor antibody, Mitochondrial ATP synthase delta subunit antibody

UniProt: P30049

Pathways: Proton Transport, Ribonucleoside Biosynthetic Process


#### **Application Details**

Application Notes: Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200,

Restrictions: For Research Use only


#### Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.




#### **Immunohistochemistry**

**Image 1.** Immunohistochemistry of paraffin-embedded human colon cancer using ABIN7144852 at dilution of 1:100



#### **Western Blotting**

Image 2. Western blot All lanes: ATP5F1D antibody at  $2\,\mu$  g/mL Lane 1: EC109 whole cell lysate Lane 2: 293T whole cell lysate Secondary Goat polyclonal to rabbit lgG at 1/15000 dilution Predicted band size: 18 kDa Observed band size: 18 kDa



#### **Immunohistochemistry**

**Image 3.** Immunohistochemistry of paraffin-embedded human lung cancer using ABIN7144852 at dilution of 1:100