
antibodies -online.com







Image



Go to Product page

_					
U	V	er	VI	е	W

Quantity:	100 μg
Target:	DHX9
Binding Specificity:	AA 325-840
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DHX9 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human ATP-dependent RNA helicase A protein (325-840AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	

Target Details

Target:	DHX9	
Alternative Name:	DHX9 (DHX9 Products)	
Background:	Background: Unwinds double-stranded DNA and RNA in a 3\' to 5\' direction. Alteration of	
	secondary structure may subsequently influence interactions with proteins or other nucleic	

acids. Functions as a transcriptional activator. Component of the CRD-mediated complex that promotes MYC mRNA stability. Involved with LARP6 in the stabilization of type I collagen mRNAs for CO1A1 and CO1A2. As component of a large PER complex is involved in the inhibition of 3\' transcriptional termination of circadian target genes such as PER1 and NR1D1 and the control of the circadian rhythms. Positively regulates HIV-1 LTR-directed gene expression.


Aliases: ATP dependent RNA helicase A antibody, ATP-dependent RNA helicase A antibody, DDX 9 antibody, DDX9 antibody, DEAD/H (Asp Glu Ala Asp/His) box polypeptide 9 antibody, DEAD/H box 9 antibody, DEAD/H box polypeptide 9 antibody, DEAH (Asp Glu Ala His) box polypeptide 9 antibody, DEAH box protein 9 antibody, DHX 9 antibody, dhx9 antibody, DHX9_HUMAN antibody, Leukophysin antibody, LKP antibody, NDH 2 antibody, NDH II antibody, NDH2 antibody, NDHII antibody, Nuclear DNA helicase II antibody, RHA antibody

UniProt:

Q08211

Application Details

Application Notes:	Recommended dilution: IHC:1:20-1:200,	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	



Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human kidney tissue using ABIN7144972 at dilution of 1:100