antibodies - online.com







anti-DDX3X antibody (AA 2-662)





Overview

Quantity:	100 μg
Target:	DDX3X
Binding Specificity:	AA 2-662
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DDX3X antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunoprecipitation (IP)

Product Details

Immunogen:	Recombinant Human ATP-dependent RNA helicase DDX3X protein (2-662AA)
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	>95%, Protein G purified

Target Details

Target:	DDX3X
Alternative Name:	DDX3X (DDX3X Products)
Background:	Background: Multifunctional ATP-dependent RNA helicase. The ATPase activity can be

stimulated by various ribo- and deoxynucleic acids indicative for a relaxed substrate specificity. In vitro can unwind partially double-stranded DNA with a preference for 5\\\'-single-stranded DNA overhangs. Is involved in several steps of gene expression, such as transcription, mRNA maturation, mRNA export and translation. However, the exact mechanisms are not known and some functions may be specific for a subset of mRNAs. Involved in transcriptional regulation. Can enhance transcription from the CDKN1A/WAF1 promoter in a SP1-dependent manner. Found associated with the E-cadherin promoter and can down-regulate transcription from the promoter. Involved in regulation of translation initiation. Proposed to be involved in positive regulation of translation such as of cyclin E1/CCNE1 mRNA and specifically of mRNAs containing complex secondary structures in their 5\\\'UTRs, these functions seem to require RNA helicase activity. Specifically promotes translation of a subset of viral and cellular mRNAs carrying a 5\\\'proximal stem-loop structure in their 5\\\'UTRs and cooperates with the eIF4F complex. Proposed to act prior to 43S ribosomal scanning and to locally destabilize these RNA structures to allow recognition of the mRNA cap or loading onto the 40S subunit. After association with 40S ribosomal subunits seems to be involved in the functional assembly of 80S ribosomes, the function seems to cover translation of mRNAs with structured and nonstructured 5\\\'UTRs and is independent of RNA helicase activity. Also proposed to inhibit capdependent translation by competetive interaction with EIF4E which can block the EIF4E:EIF4G complex formation. Proposed to be involved in stress response and stress granule assembly, the function is independent of RNA helicase activity and seems to involve association with EIF4E. May be involved in nuclear export of specific mRNAs but not in bulk mRNA export via interactions with XPO1 and NXF1. Also associates with polyadenylated mRNAs independently of NXF1. Associates with spliced mRNAs in an exon junction complex (EJC)-dependent manner and seems not to be directly involved in splicing. May be involved in nuclear mRNA export by association with DDX5 and regulating its nuclear location. Involved in innate immune signaling promoting the production of type I interferon (IFN-alpha and IFN-beta), proposed to act as viral RNA sensor, signaling intermediate and transcriptional coactivator. Involved in TBK1 and IKBKE-dependent IRF3 activation leading to IFNB induction, plays a role of scaffolding adapter that links IKBKE and IRF3 and coordinates their activation. Also found associated with IFNB promoters, the function is independent of IRF3. Can bind to viral RNAs and via association with MAVS/IPS1 and DDX58/RIG-I is thought to induce signaling in early stages of infection. Involved in regulation of apoptosis. May be required for activation of the intrinsic but inhibit activation of the extrinsic apoptotic pathway. Acts as an antiapoptotic protein through association with GSK3A/B and BIRC2 in an apoptosis antagonizing signaling complex, activation of death receptors promotes caspase-dependent cleavage of BIRC2 and DDX3X and relieves the inhibition. May be involved in mitotic chromosome segregation. Appears to be a

prime target for viral manipulations. Hepatitis B virus (HBV) polymerase and possibly vaccinia virus (VACV) protein K7 inhibit IFNB induction probably by dissociating DDX3X from TBK1 or IKBKE. Is involved in hepatitis C virus (HCV) replication, the function may involve the association with HCV core protein. HCV core protein inhibits the IPS1-dependent function in viral RNA sensing and may switch the function from a INFB inducing to a HCV replication mode. Involved in HIV-1 replication. Acts as a cofactor for XPO1-mediated nuclear export of incompletely spliced HIV-1 Rev RNAs.

Aliases: ATP dependent RNA helicase DDX3X antibody, ATP-dependent RNA helicase DDX3X antibody, CAP Rf antibody, DBX antibody, DDX14 antibody, DDX3X antibody, DDX3X_HUMAN antibody, DEAD (Asp Glu Ala Asp) box polypeptide 3 X linked antibody, DEAD (Asp-Glu-Ala-Asp) box helicase 3, X-linked antibody, DEAD box antibody, DEAD box protein 3 antibody, DEAD box protein 3 X-chromosomal antibody, DEAD box X isoform antibody, DEAD box, X isoform antibody, DEAD/H (Asp Glu Ala Asp/His) box polypeptide 3 antibody, DEAD/H box 3 antibody, DEAD/H box 3, X-linked antibody, Helicase like protein 2 antibody, Helicase-like protein 2 antibody, HLP2 antibody, X isoform antibody, X-chromosomal antibody

UniProt:

000571

Pathways:


Ribonucleoprotein Complex Subunit Organization, Positive Regulation of Endopeptidase Activity , Negative Regulation of intrinsic apoptotic Signaling, Ribosome Assembly

Application Details

Application Notes:	Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200, IF:1:50-1:200, IP:1:200-1:2000,
Restrictions:	For Research Use only


Handling

Harlanng	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.




Immunohistochemistry

Image 1. Immunohistochemistry analysis of human adrenal gland tissue using ABIN7144986 at dilution of 1:100



Western Blotting

Image 2. Western Blot Positive WB detected in: K562 whole cell lysate, Jurkat whole cell lysate, MCF-7 whole cell lysate, Rat brain tissue, Rat heart tissue, Mouse liver tissue, Mouse kidney tissue All lanes: DDX3X antibody at 3 μg/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 74, 72 kDa Observed band size: 74 kDa



Immunofluorescence

Image 3. Immunofluorescent analysis of HepG2 cells using ABIN7144986 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)

Please check the product details page for more images. Overall 4 images are available for ABIN7144986.