Datasheet for ABIN7144992 anti-DDX42 antibody (AA 519-819) (FITC)

-online.com antibodies



| $\sim$ |      |             |
|--------|------|-------------|
| ()\/#  | ervi | $\Theta(M)$ |
| 0.00   |      |             |

| Quantity:            | 100 µg                                    |
|----------------------|---|
| Target:              | DDX42                                     |
| Binding Specificity: | AA 519-819                                |
| Reactivity:          | Human                                     |
| Host:                | Rabbit                                    |
| Clonality:           | Polyclonal                                |
| Conjugate:           | This DDX42 antibody is conjugated to FITC |
| Application:         | Please inquire                            |

## Product Details

| Immunogen:        | Recombinant Human ATP-dependent RNA helicase DDX42 protein (519-819AA) |
|-------------------|--|
| Isotype:          | lgG  |
| Cross-Reactivity: | Human  |
| Purification:     | >95%, Protein G purified   |

## Target Details

| Target:           | DDX42   |  |
|-------------------|---|--|
| Alternative Name: | DDX42 (DDX42 Products)  |  |
| Background:       | Background: ATP-dependent RNA helicase. Binds to partially double-stranded RNAs (dsRNAs)      |  |
|                   | in order to unwind RNA secondary structures. Unwinding is promoted in the presence of single- |  |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7144992 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

|                     | strand binding proteins. Mediates also RNA duplex formation thereby displacing the single-    |
|---------------------|---|
|                     | strand RNA binding protein. ATP and ADP modulate its activity: ATP binding and hydrolysis by  |
|                     | DDX42 triggers RNA strand separation, whereas the ADP-bound form of the protein triggers      |
|                     | annealing of complementary RNA strands. Involved in the survival of cells by interacting with |
|                     | TP53BP2 and thereby counteracting the apoptosis-stimulating activity of TP53BP2. Relocalizes  |
|                     | TP53BP2 to the cytoplasm.   |
|                     | Aliases: DDX42ATP-dependent RNA helicase DDX42 antibody, EC 3.6.4.13 antibody, DEAD box       |
|                     | protein 42 antibody, RNA helicase-like protein antibody, RHELP antibody, RNA helicase-related |
|                     | protein antibody, RNAHP antibody, SF3b DEAD box protein antibody, Splicing factor 3B-         |
|                     | associated 125 kDa protein antibody, SF3b125 antibody   |
| UniProt:            | Q86XP3  |
| Application Details |   |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Liquid  |
| Buffer:             | Preservative: 0.03 % Proclin 300  |
|                     | Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4  |
| Preservative:       | ProClin   |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be            |
|                     | handled by trained staff only.  |
| Storage:            | -20 °C,-80 °C   |
| Storage Comment:    | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.                                 |