

Datasheet for ABIN7145068 anti-DARC antibody (AA 1-63) (FITC)



Overview

| Quantity: | 100 µg |
|----------------------|------------------------------------------|
| Target: | DARC |
| Binding Specificity: | AA 1-63 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This DARC antibody is conjugated to FITC |
| Application: | Please inquire |

Product Details

| Immunogen: | Recombinant Human Atypical chemokine receptor 1 protein (1-63AA) |
|-------------------|------------------------------------------------------------------|
| lsotype: | lgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | DARC |
|-------------------|--------------------------------------------------------------------------------------------------|
| Alternative Name: | ACKR1 (DARC Products) |
| Background: | Background: Atypical chemokine receptor that controls chemokine levels and localization via |
| | high-affinity chemokine binding that is uncoupled from classic ligand-driven signal transduction |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7145068 | 07/25/2024 | Copyright antibodies-online. All rights reserved. cascades, resulting instead in chemokine sequestration, degradation, or transcytosis. Also known as interceptor (internalizing receptor) or chemokine-scavenging receptor or chemokine decoy receptor. Has a promiscuous chemokine-binding profile, interacting with inflammatory chemokines of both the CXC and the CC subfamilies but not with homeostatic chemokines. Acts as a receptor for chemokines including CCL2, CCL5, CCL7, CCL11, CCL13, CCL14, CCL17, CXCL5, CXCL6, IL8/CXCL8, CXCL11, GR0, RANTES, MCP-1, TARC and also for the malaria parasites P.vivax and P.knowlesi. May regulate chemokine bioavailability and, consequently, leukocyte recruitment through two distinct mechanisms: when expressed in endothelial cells, it sustains the abluminal to luminal transcytosis of tissue-derived chemokines and their subsequent presentation to circulating leukocytes, when expressed in erythrocytes, serves as blood reservoir of cognate chemokines but also as a chemokine sink, buffering potential surges in plasma chemokine levels.

Aliases: ACKR1 antibody, Atypical chemokine receptor 1 antibody, CCBP1 antibody, CD234 antibody, DARC antibody, Dfy antibody, Duffy antigen/chemokine receptor antibody, duffy blood group antibody, Duffy blood group antigen antibody, Duffy blood group chemokine receptor antibody, DUFFY_HUMAN antibody, FY antibody, Fy glycoprotein antibody, Glycoprotein D antibody, GPD antibody, GpFy antibody, Plasmodium vivax receptor antibody, WBCQ1 antibody

UniProt:

Q16570

Application Details

Restrictions: For Research Use only Handling Format: Liquid Buffer: Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4 ProClin Preservative: Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. -20 °C.-80 °C Storage: Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN7145068 | 07/25/2024 | Copyright antibodies-online. All rights reserved.