antibodies -online.com





anti-BBS2 antibody (AA 1-96)

3 Images



Go to Product page

Overview

Quantity:	100 μL
Target:	BBS2
Binding Specificity:	AA 1-96
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BBS2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Bardet-Biedl syndrome 2 protein (1-96AA)
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Purification:	Antigen Affinity Purified

Target Details

Target:	BBS2
Alternative Name:	BBS2 (BBS2 Products)
Background:	Background: The BBSome complex is thought to function as a coat complex required for
	sorting of specific membrane proteins to the primary cilia. The BBSome complex is required for

ciliogenesis but is dispensable for centriolar satellite function. This ciliogenic function is mediated in part by the Rab8 GDP/GTP exchange factor, which localizes to the basal body and contacts the BBSome. Rab8(GTP) enters the primary cilium and promotes extension of the ciliary membrane. Firstly the BBSome associates with the ciliary membrane and binds to RAB3IP/Rabin8, the guanosyl exchange factor (GEF) for Rab8 and then the Rab8-GTP localizes to the cilium and promotes docking and fusion of carrier vesicles to the base of the ciliary membrane. The BBSome complex, together with the LTZL1, controls SMO ciliary trafficking and contributes to the sonic hedgehog (SHH) pathway regulation. Required for proper BBSome complex assembly and its ciliary localization.

Aliases: BBS2Bardet-Biedl syndrome 2 protein antibody

UniProt: Q9BXC9

Pathways: Hedgehog Signaling


Application Details

Application Notes: Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200,

Restrictions: For Research Use only


Handling

Format:	Liquid
Buffer:	PBS with 0.02 % sodium azide, 50 % glycerol, pH 7.3.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.




Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human endometrial cancer using ABIN7145239 at dilution of 1:100



Western Blotting

Image 2. Western blot All lanes: BBS2 antibody at $5.01 \,\mu$ g/mL Lane 1: Mouse liver tissue Lane 2: Mouse brain tissue Lane 3: Mouse gonadal tissue Secondary Goat polyclonal to rabbit IgG at 1/10000 dilution Predicted band size: 80 kDa Observed band size: 80 kDa



Immunohistochemistry

Image 3. Immunohistochemistry of paraffin-embedded human testis tissue using ABIN7145239 at dilution of 1:100