Datasheet for ABIN7145278

anti-BATF antibody (AA 1-125) (HRP)

Go to Product page

Overview

Quantity:	100 µg
Target:	BATF
Binding Specificity:	AA 1-125
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BATF antibody is conjugated to HRP
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Basic leucine zipper transcriptional factor ATF-like protein (1-125AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	BATF
Alternative Name:	BATF (BATF Products)
Background:	Background: AP-1 family transcription factor that controls the differentiation of lineage-specific cells in the immune system: specifically mediates the differentiation of T-helper 17 cells (Th17),

Target Details

	follicular T-helper cells (TfH), CD8(+) dendritic cells and class-switch recombination (CSR) in B-cells. Acts via the formation of a heterodimer with JUNB that recognizes and binds DNA sequence 5\'-TGA[CG]TCA-3\'. The BATF-JUNB heterodimer also forms a complex with IRF4 (or IRF8) in immune cells, leading to recognition of AICE sequence (5\'-TGAnTCA/GAAA-3\'), an immune-specific regulatory element, followed by cooperative binding of BATF and IRF4 (or IRF8) and activation of genes. Controls differentiation of T-helper cells producing interleukin-17 (Th17 cells) by binding to Th17-associated gene promoters: regulates expression of the transcription factor RORC itself and RORC target genes such as IL17 (IL17A or IL17B). Also involved in differentiation of follicular T-helper cells (TfH) by directing expression of BCL6 and MAF. In B-cells, involved in class-switch recombination (CSR) by controlling the expression of both AICDA and of germline transcripts of the intervening heavy-chain region and constant heavy-chain region (I(H)-C(H)). Following infection, can participate in CD8(+) dendritic cell differentiation via interaction with IRF4 and IRF8 to mediate cooperative gene activation. Regulates effector CD8(+) T-cell differentiation by regulating expression of SIRT1. Following DNA damage, part of a differentiation checkpoint that limits self-renewal of hematopoietic stem cells (HSCs): up-regulated by STAT3, leading to differentiation of HSCs, thereby restricting self-renewal of HSCs (By similarity). Aliases: Activating transcription factor B antibody, B ATF antibody, B-ATF antibody, B-cell-activating transcription factor antibody, Basic leucine zipper transcription factor like antibody, Basic leucine zipper transcriptional factor ATF like antibody, Basic leucine zipper transcriptional factor ATF-like antibody, Batf antibody, BATF_HUMAN antibody, BATF1 antibody, SF HT activated gene 2 protein antibody, SF-HT-activated gene 2 protein antibody, SFA 2 antibody, SFA-2 antibody, SFA2 antibody
UniProt:	Q16520

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

 Copyright antibodies-online. All rights reserved.