

[Go to Product page](#)

Datasheet for ABIN7145315

**anti-BOK antibody (AA 15-103) (Biotin)**

## Overview

Quantity:	100 µg
Target:	BOK
Binding Specificity:	AA 15-103
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BOK antibody is conjugated to Biotin
Application:	ELISA

## Product Details

Immunogen:	Recombinant Human Bcl-2-related ovarian killer protein (15-103AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	BOK
Alternative Name:	BOK ( <a href="#">BOK Products</a> )
Background:	Background: Isoform 1: Apoptosis regulator that functions through different apoptotic signaling pathways (PubMed:27076518, PubMed:15102863, PubMed:20673843). Plays a roles as pro-

## Target Details

apoptotic protein that positively regulates intrinsic apoptotic process in a BAX- and BAK1-dependent manner or in a BAX- and BAK1-independent manner (PubMed:27076518, PubMed:15102863). In response to endoplasmic reticulum stress promotes mitochondrial apoptosis through downstream BAX/BAK1 activation and positive regulation of PERK-mediated unfolded protein response (By similarity). Activates apoptosis independently of heterodimerization with survival-promoting BCL2 and BCL2L1 through induction of mitochondrial outer membrane permeabilization, in a BAX- and BAK1-independent manner, in response to inhibition of ERAD-proteasome degradation system, resulting in cytochrome c release (PubMed:27076518). In response to DNA damage, mediates intrinsic apoptotic process in a TP53-dependent manner (PubMed:15102863). Plays a role in granulosa cell apoptosis by CASP3 activation (PubMed:20673843). Plays a roles as anti-apoptotic protein during neuronal apoptotic process, by negatively regulating poly ADP-ribose polymerase-dependent cell death through regulation of neuronal calcium homeostasis and mitochondrial bioenergetics in response to NMDA excitation (By similarity). In addition to its role in apoptosis, may regulate trophoblast cell proliferation during the early stages of placental development, by acting on G1/S transition through regulation of CCNE1 expression (PubMed:19942931). May also play a role as an inducer of autophagy by disrupting interaction between MCL1 and BECN1 (PubMed:24113155).

Aliases: Bcl 2 related ovarian killer protein antibody, Bcl-2-like protein 9 antibody, Bcl-2-related ovarian killer protein antibody, BCL2 related ovarian killer antibody, Bcl2-L-9 antibody, BCL2L9 antibody, BOK antibody, BOK\_HUMAN antibody, BOKL antibody, Hbok antibody, MGC4631 antibody

UniProt: [Q9UMX3](#)

Pathways: [Positive Regulation of Endopeptidase Activity](#)

## Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

## Handling

---

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.