antibodies - online.com







anti-SNTB1 antibody (AA 51-111)





\sim	
Over\	$/1 \triangle 1 \Lambda /$
OVEIV	/ I C V V

Quantity:	100 μg
Target:	SNTB1
Binding Specificity:	AA 51-111
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SNTB1 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant Human Beta-1-syntrophin protein (51-111AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	SNTB1
Alternative Name:	SNTB1 (SNTB1 Products)
Background:	Background: Adapter protein that binds to and probably organizes the subcellular localization of a variety of membrane proteins. May link various receptors to the actin cytoskeleton and the

dystrophin glycoprotein complex.

Aliases: 59 DAP antibody, 59 kDa dystrophin associated protein A1 basic component 1 antibody, 59 kDa dystrophin-associated protein A1 basic component 1 antibody, A1B antibody, Beta 1 syntrophin antibody, Beta-1-syntrophin antibody, BSYN2 antibody, DAPA1B antibody, Dystrophin associated protein A1 59 kDa basic component 1 antibody, FLJ22442 antibody, MGC111389 antibody, SNT2 antibody, SNT2B1 antibody, SNTB 1 antibody, Sntb1 antibody, SNTB1_HUMAN antibody, Syntrophin 2 antibody, Syntrophin beta 1 antibody, Syntrophin beta 1 (dystrophin associated protein A1 59 kDa basic component 1) antibody, Syntrophin-2 antibody, Tax interaction protein 43 antibody, TIP 43 antibody, TIP-43 antibody

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be

UniProt:

Q13884

Application Details

Precaution of Use:


Storage:

Application Notes:	Recommended dilution: IF:1:50-1:200,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Format: Buffer:	Liquid Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

-20 °C,-80 °C

handled by trained staff only.



Immunofluorescence

Image 1. Immunofluorescence staining of HepG2 cells with ABIN7145359 at 1:66, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).