# antibodies -online.com





## anti-FGF22 antibody (AA 23-100) (Cy3)



Go to Product page

| ( ) | 11/               | IN   | /ie    | A .    |
|-----|-------------------|------|--------|--------|
|     | / // <del> </del> | ۱ ات | /   (− | ' \/\/ |
|     |                   |      |        |        |

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | FGF22  |
| Binding Specificity: | AA 23-100  |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This FGF22 antibody is conjugated to Cy3   |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human FGF22. |  |
|-----------------------|--|--|
| Isotype:              | IgG  |  |
| Predicted Reactivity: | Human,Mouse,Cow  |  |
| Purification:         | Purified by Protein A.                                     |  |

#### **Target Details**

| Target:           | FGF22  |
|-------------------|--|
| Alternative Name: | FGF22 (FGF22 Products)   |
| Background:       | Synonyms: FGF 22, FGF-22, FGF22, FGF22_HUMAN, Fibroblast growth factor 22, si:dkey |

| 01 | -0   | $\circ$ |
|----|------|---------|
|    | lez. | . Z     |

Background: FGF22 is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities and are involved in a variety of biological processes including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. The mouse homolog of this gene was found to be preferentially expressed in the inner root sheath of the hair follicle, which suggested a role in hair development. Fibroblast Growth Factor-22 stimulates the proliferation and activation of cells that express FGF receptor.

Gene ID: 27006

Pathways: RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin

Signaling Pathway

### **Application Details**

| Application Notes: | IF(IHC-P) 1:50-200 |
|--------------------|--------------------|
|                    | IF(IHC-F) 1:50-200 |
|                    | IF(ICC) 1:50-200   |

Restrictions: For Research Use only

#### Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |