# antibodies .- online.com







## anti-CHN2 antibody (AA 1-91)



Image



#### Overview

Quantity:	100 μg
Target:	CHN2
Binding Specificity:	AA 1-91
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CHN2 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

#### **Product Details**

Immunogen:	Recombinant Human Beta-chimaerin protein (1-91AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	CHN2
Alternative Name:	CHN2 (CHN2 Products)
Background:	Background: GTPase-activating protein for p21-rac. Insufficient expression of beta-2 chimaerin
	is expected to lead to higher Rac activity and could therefore play a role in the progression from

#### Target Details

	low-grade to high-grade tumors.
	Aliases: CHN2 antibody, ARHGAP3 antibody, BCH antibody, Beta-chimaerin antibody, Beta-
	chimerin antibody, Rho GTPase-activating protein 3 antibody
UniProt:	P52757


#### **Application Details**

Application Notes:	Recommended dilution: IHC:1:500-1:1000,
Restrictions:	For Research Use only

#### Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

#### **Images**



#### **Immunohistochemistry**

**Image 1.** IHC image of ABIN7145458 diluted at 1:800 and staining in paraffin-embedded human liver tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.