antibodies -online.com





anti-EPHX2 antibody (AA 1-555) (Biotin)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μg
Target:	EPHX2
Binding Specificity:	AA 1-555
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EPHX2 antibody is conjugated to Biotin
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Bifunctional epoxide hydrolase 2 protein (1-555AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	

Target Details

Target:	EPHX2
Alternative Name:	EPHX2 (EPHX2 Products)
Background:	Background: Bifunctional enzyme. The C-terminal domain has epoxide hydrolase activity and
	acts on epoxides (alkene oxides, oxiranes) and arene oxides. Plays a role in xenobiotic

Target Details

metabolism by degrading potentially toxic epoxides. Also determines steady-state levels of physiological mediators. The N-terminal domain has lipid phosphatase activity, with the highest activity towards threo-9,10-phosphonooxy-hydroxy-octadecanoic acid, followed by erythro-9,10-phosphonooxy-hydroxy-octadecanoic acid, 12-phosphonooxy-octadec-9Z-enoic acid, 12-phosphonooxy-octadec-9E-enoic acid, and p-nitrophenyl phospate.

Aliases: Bifunctional epoxide hydrolase 2 antibody, CEH antibody, Cytosolic epoxide hydrolase antibody, EPHX2 antibody, Epoxide hydrolase 2 antibody, Epoxide hydrolase 2 antibody, Epoxide hydrolase 2 cytoplasmic antibody, epoxide hydrolase 2, cytosolic antibody, Epoxide hydrolase soluble antibody, HYES_HUMAN antibody, SEH antibody, Soluble epoxide hydrolase antibody

UniProt:

P34913

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	