antibodies -online.com







anti-EPHX2 antibody (AA 1-555) (FITC)



Go to Product page

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μg
Target:	EPHX2
Binding Specificity:	AA 1-555
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EPHX2 antibody is conjugated to FITC
Application:	Please inquire

Product Details

Immunogen:	Recombinant Human Bifunctional epoxide hydrolase 2 protein (1-555AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	EPHX2	
Alternative Name:	EPHX2 (EPHX2 Products)	
Background:	Background: Bifunctional enzyme. The C-terminal domain has epoxide hydrolase activity at acts on epoxides (alkene oxides, oxiranes) and arene oxides. Plays a role in xenobiotic	

Target Details

metabolism by degrading potentially toxic epoxides. Also determines steady-state levels of physiological mediators. The N-terminal domain has lipid phosphatase activity, with the highest activity towards threo-9,10-phosphonooxy-hydroxy-octadecanoic acid, followed by erythro-9,10-phosphonooxy-hydroxy-octadecanoic acid, 12-phosphonooxy-octadec-9Z-enoic acid, 12-phosphonooxy-octadec-9E-enoic acid, and p-nitrophenyl phospate.

Aliases: Bifunctional epoxide hydrolase 2 antibody, CEH antibody, Cytosolic epoxide hydrolase antibody, EPHX2 antibody, Epoxide hydratase antibody, Epoxide hydrolase 2 antibody, Epoxide hydrolase 2 cytoplasmic antibody, epoxide hydrolase 2, cytosolic antibody, Epoxide hydrolase soluble antibody, HYES_HUMAN antibody, SEH antibody, Soluble epoxide hydrolase antibody

UniProt:

P34913

Application Details

Restrictions:

For Research Use only

Handling

Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	