antibodies .- online.com







anti-NR1H4 antibody (AA 1-476) (FITC)



()	11/0	K\ /	iew	1
	\cup	ועוי	$\square \vee \vee$	ı

Quantity:	100 μg
Target:	NR1H4
Binding Specificity:	AA 1-476
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NR1H4 antibody is conjugated to FITC
Application:	Please inquire

Product Details

Immunogen:	Recombinant Human Bile acid receptor protein (1-476AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	

Target Details

Target:	NR1H4
Alternative Name:	NR1H4 (NR1H4 Products)
Background:	Background: Ligand-activated transcription factor. Receptor for bile acids such as
	chenodeoxycholic acid, lithocholic acid and deoxycholic acid. Represses the transcription of the

cholesterol 7-alpha-hydroxylase gene (CYP7A1) through the induction of NR0B2 or FGF19 expression, via two distinct mechanisms. Activates the intestinal bile acid-binding protein (IBABP). Activates the transcription of bile salt export pump ABCB11 by directly recruiting histone methyltransferase CARM1 to this locus.

Aliases: BAR antibody, Bile acid receptor antibody, Farnesoid X activated receptor antibody, Farnesoid X receptor antibody, Farnesoid X-activated receptor antibody, Farnesol receptor HRR 1 antibody, Farnesol receptor HRR-1 antibody, Farnesol receptor HRR1 antibody, FXR antibody, HRR 1 antibody, HRR1 antibody, NR1H4 antibody, NR1H4_HUMAN antibody, Nuclear receptor subfamily 1 group H member 4 antibody, Retinoid X receptor interacting protein 14 antibody, Retinoid X receptor-interacting protein 14 antibody, RIP 14 antibody, RIP14 antibody, RXR interacting protein 14 antibody, RXR-interacting protein 14 antibody

UniProt:

096RI1

Pathways:

Nuclear Receptor Transcription Pathway, Steroid Hormone Mediated Signaling Pathway, Regulation of Carbohydrate Metabolic Process

Application Details

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.